



## EDAPT INFORMATION FOR SCHOOLS

### What is EDAPT?

The Early Diagnosis And Preventative Treatment of Psychotic Illness (EDAPT) provides confidential assessment and early assistance for young people between the ages of 12 and 25 in the Greater Sacramento Area who are at risk for mental illnesses. Most services provided by EDAPT staff are funded through research grants. Contact EDAPT for an initial assessment or consultation at no charge.

By getting help early, a person's chances greatly improve for staying in school, working, maintaining friendships and planning for the future. The EDAPT Program provides education and treatment for young people and their families, and conducts research about the effects of early identification in stemming the progression of mental illness.

### How can I tell the difference between the early warning signs for psychosis and other mental health or developmental issues?

Early warning signs can appear to be laziness, forgetfulness, clinical depression or Attention Deficit Hyperactivity Disorder (ADHD). The "red flags" for psychosis include social withdrawal, a drop in functioning, peculiar behavior, heightened sensory sensitivity, altered perceptions, loss of motivation, dramatic sleep and appetite changes, suspiciousness of others and unusual or exaggerated beliefs. These symptoms in isolation and for shorter lengths of time could be a response to a loss or significant change in one's life. A combination and worsening of these symptoms could preface the onset of psychosis. If a professional or parent observes some of these changes in a young person, we encourage them to contact us for an assessment.

### What parts of the brain are affected in young people with prodromal psychosis?

The frontal lobes of the human brain are the area of higher order cognitive functions like judgment, planning, reasoning, organization, maintaining a focus and initiating action. During adolescence, maturation of the frontal lobes is incomplete. It is during this stage of development that frontal lobe impairment may become evident in a young person at risk of a major mental illness. This under activity in the frontal lobes can contribute to a clinical picture of diminishing academic performance, trouble processing new information, disorganization and apathy or a lack of mental liveliness. A corresponding over activity of the limbic system impacts primitive emotions and feelings and creates a sensory overload experience which can trigger psychosis.

### How can EDAPT be helpful to schools?

EDAPT offers a range of services to schools in the Greater Sacramento Area and beyond, including professional training on understanding psychosis and recognizing the early symptoms, consultation on how to work with a child who may be at risk and information about how and when to make a referral to EDAPT. We are also available to present to student groups (such as health education classes or school-wide assemblies). These interactive presentations help to reduce stigma and misinformation about mental illness, as well as provide students with information about early treatment.

EDAPT also provides an educational toolkit for teachers who are interested in adding mental health content to their curricula. Our training and materials may be helpful for school staff to reduce stigma and support students who may be struggling with mental health concerns or symptoms.

